


# HOW TO TAKE OXBRYTA® (voxelotor) 300 MG TABLETS FOR ORAL SUSPENSION

FOR PATIENTS 4 YEARS AND  
OLDER BASED ON WEIGHT AND  
ABILITY TO SWALLOW TABLETS



TULLY & HAJAR ... Sisters, artists, and actual Oxbryta patients

## WHAT YOU'LL NEED



- A teaspoon
- A small cup
- A room temperature clear drink  
(For example: drinking water, clear soda,  
apple juice, clear electrolyte drinks, clear  
flavored drinks, or clear sports drinks)
- The prescribed number of Oxbryta  
300 mg Tablets for Oral Suspension  
(light yellow, round shaped, and marked  
with "300 D" on one side)



For complete information on  
Dosage and Administration,  
see the detailed [Instructions  
for Use on Oxbryta.com](#).



## WATCH NOW

Scan the QR code to see how a  
healthcare provider prepares Oxbryta  
300 mg Tablets for Oral Suspension.

## IMPORTANT INFORMATION

- Take Oxbryta exactly as your healthcare provider tells you
- **Do not** change your dose or stop taking Oxbryta unless your healthcare provider tells you to. Your healthcare provider may change your dose if needed
- Take your prescribed dose of Oxbryta 1 time each day
- Take Oxbryta with or without food
- **Do not** swallow whole, cut, crush, or chew the tablets for oral suspension
- **Check to make sure you receive the correct dosage form of Oxbryta each time your prescription is filled to avoid taking the wrong medicine.** Contact your pharmacist or healthcare provider if you did not receive the correct medication
- Your healthcare provider may also prescribe a medicine called hydroxyurea during treatment with Oxbryta
- **If you or your child miss a dose or do not take the entire dose of Oxbryta,** skip that dose and return to your normal dosing schedule the next day
- Do not take St. John's wort during treatment with Oxbryta

## INDICATION

### What is OXBRYTA?

OXBRYTA is a prescription medicine used for the treatment of sickle cell disease in adults and children 4 years of age and older. It is not known if OXBRYTA is safe and effective in children with sickle cell disease below 4 years of age.

OXBRYTA is approved under accelerated approval based on an increase in hemoglobin. Continued approval may depend on verification of clinical benefits in ongoing trials.

**Please see [Important Safety Information](#) on page 4 and [Patient Information](#) on [Oxbryta.com](#).**

**Oxbryta**  
(voxelotor)  
300mg • 500mg tablets  
300mg tablets for oral suspension

# STEPS FOR PREPARING AND TAKING OR GIVING OXBRYTA 300 MG TABLETS FOR ORAL SUSPENSION

1



## POUR CLEAR DRINK

Be sure your hands are clean and dry, then pour the correct amount of room temperature clear drink in the cup.

- The table on the right shows the amount of clear drink needed for the prescribed dose
- You may add more clear drink if needed to mix the tablets
- Your healthcare provider will tell you how many Oxbryta 300 mg Tablets for Oral Suspension you will need for your or your child's dose

NUMBER OF OXBRYTA  
300 MG TABLETS FOR  
ORAL SUSPENSION

AMOUNT OF  
CLEAR DRINK

1	1 tsp (5 mL)
2	2 tsp (10 mL)
3	3 tsp (15 mL)
4	4 tsp (20 mL)
5	5 tsp (25 mL)
7	7 tsp (35 mL)
8	8 tsp (40 mL)

2



## ADD PRESCRIBED NUMBER OF TABLETS

Visually inspect the pills to ensure they are the correct medication. Please see photo to the right.

- Add the prescribed number of Oxbryta 300 mg Tablets for Oral Suspension to the cup
- **Do not** swallow whole, cut, crush, or chew the tablets for oral suspension



Not actual size

3




## SWIRL & WAIT

Swirl the cup until the tablet(s) breaks apart in the drink. Be careful not to spill. The tablet(s) will not completely dissolve. You will still see small tablet clumps in the mixture.

- If you spill any medicine, clean up the spill. Throw away the rest of the prepared medicine and make a new dose

Wait for 1 to 5 minutes before taking or giving the dose.



Please see **Important Safety Information** on page 4 and **Patient Information** on [Oxbryta.com](http://Oxbryta.com).

4




## SWIRL & DRINK

Swirl the cup again. Be careful not to spill the mixture. Drink all the prepared medicine right away.

- If giving to a child, make sure that the child is upright. Give all the prepared medicine to the child right away



5



## ADD MORE CLEAR DRINK & REPEAT

Add 1 or 2 more teaspoons of room temperature clear drink to the cup. This is to mix the leftover medicine and to make sure the full dose is taken. Swirl the cup until all the leftover medicine is mixed, and drink it right away. Repeat this step until all the medicine is taken.

- After taking all the medicine, you or your child may drink more water or any type of drink
- Be sure to wash the spoon and cup with warm water and soap



## HOW TO STORE OXBRYTA

- Store Oxbryta at 68°F to 86°F (20°C to 30°C)
  - Oxbryta comes in a child-resistant package
  - The bottle contains a polyester coil. **Do not** eat the polyester coil
- Keep Oxbryta and all medicines out of the reach of children.**

For full information on how you should take Oxbryta, please see the [Patient Information on Oxbryta.com](https://www.oxbryta.com). If you have any questions, please talk to your healthcare provider.


**Oxbryta**<sup>®</sup>  
 (voxelotor)  
 300mg • 500mg tablets  
 300mg tablets for oral suspension

## INDICATION AND IMPORTANT SAFETY INFORMATION

### INDICATION

#### What is OXBRYTA?

OXBRYTA is a prescription medicine used for the treatment of sickle cell disease in adults and children 4 years of age and older. It is not known if OXBRYTA is safe and effective in children with sickle cell disease below 4 years of age.

OXBRYTA is approved under accelerated approval based on an increase in hemoglobin. Continued approval may depend on verification of clinical benefits in ongoing trials.

### IMPORTANT SAFETY INFORMATION

**Do not take OXBRYTA** if you or your child have had an allergic reaction to voxelotor or any of the ingredients in OXBRYTA. See the end of the patient leaflet for a list of the ingredients in OXBRYTA.

**Before taking OXBRYTA, tell your healthcare provider about all of your medical conditions, including if you or your child:**

- have liver problems
- are pregnant or plan to become pregnant. It is not known if OXBRYTA can harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if OXBRYTA can pass into your breastmilk and if it can harm your baby. Do not breastfeed during treatment with OXBRYTA and for at least 2 weeks after the last dose.

**Tell your healthcare provider about all the medicines you or your child take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Some medicines may affect how OXBRYTA works. OXBRYTA may also affect how other medicines work and may affect the results of certain blood tests. Keep a list of all your medicines and show it to your healthcare provider.

#### What should I avoid while taking OXBRYTA?

Do not take St. John's wort during treatment with OXBRYTA.

#### What are the possible side effects of OXBRYTA?

**OXBRYTA can cause serious side effects, including:**

- **Severe skin rash and serious allergic reactions.** Treatment with OXBRYTA may cause severe skin reactions and serious allergic reactions. The organs in your body may also be affected, such as your liver, kidneys or lungs, and your blood cells.
- **Stop taking OXBRYTA**, and tell your healthcare provider or get emergency medical help right away if you develop any of the following signs or symptoms during treatment:
  - rash
  - hives
  - high temperature (fever)
  - swollen glands (lymph nodes)
  - trouble swallowing
  - shortness of breath (difficult breathing)
  - swelling of your face, around your eyes, lips, or tongue
  - lack of energy and tiredness (fatigue)
  - muscle or joint aches

**The most common side effects of OXBRYTA include:**

- headache
- nausea or vomiting
- fever
- stomach-area (abdominal) pain
- diarrhea
- rash or hives

These are not all the possible side effects of OXBRYTA.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Pfizer Inc. at 1-800-438-1985.

**Keep OXBRYTA and all medicines out of the reach of children.**

**Please see [Patient Information on Oxbryta.com](#).**



Oxbryta® is a registered trademark of Global Blood Therapeutics, Inc., a wholly owned subsidiary of Pfizer Inc.  
PP-LTV-USA-2556 © 2023 Pfizer Inc. All rights reserved. November 2023

  
300mg • 500mg tablets  
300mg tablets for oral suspension